

Subject:	Anesthesia Services and Moderate (“Conscious”) Sedation	Publish Date:	12/16/2020
Guideline #:	CG-MED-21	Last Review Date:	11/05/2020
Status:	Reviewed		

Description

This document addresses the medical necessity of anesthesia services. Anesthesia services include all services associated with the administration and monitoring of analgesia or anesthesia in order to produce partial or complete loss of sensation. Examples of various methods of anesthesia include general anesthesia, regional anesthesia, monitored anesthesia care (MAC), moderate sedation (“conscious sedation”), and local infiltration or topical application. This document does not address anesthesia services performed during gastrointestinal endoscopic procedures.

Note: Please see the following documents for additional information:

- CG-MED-34 Monitored Anesthesia Care for Gastrointestinal Endoscopic Procedures
- CG-MED-41 Moderate to Deep Anesthesia Services for Dental Surgery in the Facility Setting
- CG-MED-65 Manipulation Under Anesthesia
- CG-MED-78 Anesthesia Services for Interventional Pain Management Procedures

Note: This document does not address whether or not reimbursement is provided for the anesthesia service and is **not** intended to explain the billing and reimbursement of anesthesia.

Clinical Indications

Medically Necessary:

General Anesthesia or Regional Anesthesia

Administration of general or regional anesthesia is considered **medically necessary** when both of the following criteria are met:

- A. The services are provided by an individual other than the attending physician performing the procedure;
and
- B. Alternative types of anesthesia, sedation, or analgesia are not appropriate.

If general or regional anesthesia is requested for a procedure typically not requiring either of these levels of anesthesia service, a medical necessity review will be performed. This review will assess not only the procedure involved, but also other individual-specific issues, such as age, mental status, ability to cooperate, co-morbid conditions, and general medical status.

Monitored Anesthesia Care (MAC)

Monitored anesthesia care (MAC) is considered **medically necessary** when **all** of the following criteria are met:

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue’s standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue’s Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Anesthesia Services and Moderate (“Conscious”) Sedation

- A. MAC is requested by the attending physician; **and**
- B. The services are provided by an individual other than the attending physician performing the procedure; **and**
- C. Qualified anesthesia personnel (anesthesiologists or qualified anesthetists such as certified registered nurse anesthetists) administering monitored anesthesia care are continuously present to monitor the individual and provide anesthesia care; **and**
- D. The individual's medical condition requires medical direction or supervision of the anesthetic to ensure control of the sedation, medication, and airway, and to prevent sudden changes in condition from disrupting the procedure and placing the individual at risk; **and**
- E. Constant monitoring of the individual's vital signs is provided to anticipate the need for general anesthesia administration or for the treatment of adverse physiologic reactions such as hypotension, excessive pain, difficulty breathing, arrhythmias, adverse drug reactions, etc. In addition, the possibility that the procedure may become more extensive, or result in unforeseen complications, requires comprehensive monitoring or anesthetic intervention; **and**
- F. Appropriate documentation is available to reflect pre- and post-anesthetic evaluations and intraoperative monitoring.

Anesthesia Services including MAC for Surgical Procedures

For surgical procedures which do not usually require anesthesia services, anesthesia services including monitored anesthesia care (MAC) are considered **medically necessary** when the individual's condition requires the presence of qualified anesthesia personnel to perform monitored anesthesia in addition to the physician performing the procedure, and is so documented. The medical condition must be significant enough to impact the need to provide anesthesia services including MAC. Complex procedures and procedures in high-risk individuals may justify the use of an anesthesiologist or anesthetist to provide conscious sedation or deep sedation. See Appendix for physical status classifications. The presence of a stable, treated condition of itself is not necessarily sufficient.

Moderate (“Conscious”) Sedation

Moderate sedation (“conscious sedation”) ordered by the attending physician and administered by the surgeon or physician performing the procedure or an independent trained practitioner is considered **medically necessary** when alternative types of anesthesia, sedation, or analgesia are not appropriate.

Local Anesthesia

The administration of local anesthesia is considered **medically necessary** when alternative types of anesthesia, sedation, or analgesia are not appropriate.

Standby Anesthesia Services

Standby anesthesia service is when the anesthesiologist would be immediately available if a clinical need should arise but the anesthesiologist may be elsewhere performing other duties. Stand-by anesthesia is considered **medically necessary** when a procedure, which does not normally require anesthesia services, has a significant potential for catastrophic complications or potential for the need of other intervention that would require immediate availability of general anesthesia.

Not Medically Necessary:

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Anesthesia Services and Moderate (“Conscious”) Sedation

Anesthesia services are considered **not medically necessary** for all other indications.

Coding

The following codes for treatments and procedures applicable to this document are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

When services may be Medically Necessary when criteria are met:

CPT

00100-00222	Anesthesia for procedures on the head [includes codes 00100, 00102, 00103, 00104, 00120, 00124, 00126, 00140, 00142, 00144, 00145, 00147, 00148, 00160, 00162, 00164, 00170, 00172, 00174, 00176, 00190, 00192, 00210, 00211, 00212, 00214, 00215, 00216, 00218, 00220, 00222]
00300-00352	Anesthesia for procedures on the neck [includes codes 00300, 00320, 00322, 00326, 00350, 00352]
00400-00474	Anesthesia for procedures on the thorax [includes codes 00400, 00402, 00404, 00406, 00410, 00450, 00454, 00470, 00472, 00474]
00500-00580	Anesthesia for intrathoracic procedures [includes codes 00500, 00520, 00522, 00524, 00528, 00529, 00530, 00532, 00534, 00537, 00539, 00540, 00541, 00542, 00546, 00548, 00550, 00560, 00561, 00562, 00563, 00566, 00567, 00580]
00600-00670	Anesthesia for procedures on spine and spinal cord [includes codes 00600, 00604, 00620, 00625, 00626, 00630, 00632, 00635, 00640, 00670]
00700-00797	Anesthesia for procedures on upper abdomen [includes codes 00700, 00702, 00730, 00750, 00752, 00754, 00756, 00770, 00790, 00792, 00794, 00796, 00797]
00800-00882	Anesthesia for procedures on lower abdomen [includes codes 00800, 00802, 00820, 00830, 00832, 00834, 00836, 00840, 00842, 00844, 00846, 00848, 00851, 00860, 00862, 00864, 00865, 00866, 00868, 00870, 00872, 00873, 00880, 00882]
00902-00952	Anesthesia for procedures on perineum [includes codes 00902, 00904, 00906, 00908, 00910, 00912, 00914, 00916, 00918, 00920, 00921, 00922, 00924, 00926, 00928, 00930, 00932, 00934, 00936, 00938, 00940, 00942, 00944, 00948, 00950, 00952]
01112-01173	Anesthesia for procedures on pelvis [includes codes 01112, 01120, 01130, 01140, 01150, 01160, 01170, 01173]
01200-01274	Anesthesia for procedures on upper leg [includes codes 01200, 01202, 01210, 01212, 01214, 01215, 01220, 01230, 01232, 01234, 01250, 01260, 01270, 01272, 01274]
01320-01444	Anesthesia for procedures on knee and popliteal area [includes codes 01320, 01340, 01360, 01380, 01382, 01390, 01392, 01400, 01402, 01404, 01420, 01430, 01432, 01440, 01442, 01444]
01462-01522	Anesthesia for procedures on lower leg [includes codes 01462, 01464, 01470, 01472, 01474, 01480, 01482, 01484, 01486, 01490, 01500, 01502, 01520, 01522]
01610-01680	Anesthesia for procedures on shoulder and axilla [includes codes 01610, 01620, 01622, 01630, 01634, 01636, 01638, 01650, 01652, 01654, 01656, 01670, 01680]

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Anesthesia Services and Moderate (“Conscious”) Sedation

01710-01782	Anesthesia for procedures on upper arm and elbow [includes codes 01710, 01712, 01714, 01716, 01730, 01732, 01740, 01742, 01744, 01756, 01758, 01760, 01770, 01772, 01780, 01782]
01810-01860	Anesthesia for procedures on forearm, wrist, and hand [includes codes 01810, 01820, 01829, 01830, 01832, 01840, 01842, 01844, 01850, 01852, 01860]
01916-01933	Anesthesia for radiological procedures [includes codes 01916, 01920, 01922, 01924, 01925, 01926, 01930, 01931, 01932, 01933]
01951-01953	Anesthesia for second- and third-degree burn excision or debridement with or without skin grafting, any site, for total body surface area (TBSA) treated during anesthesia and surgery [includes codes 01951, 01952, 01953]
01958-01969	Anesthesia for obstetric procedures [includes codes 01958, 01960, 01961, 01962, 01963, 01965, 01966, 01967, 01968, 01969]
01990	Physiological support for harvesting of organ(s) from brain-dead patient
01996	Daily hospital management of epidural or subarachnoid continuous drug administration
01999	Unlisted anesthesia procedure(s)
99100	Anesthesia for patient of extreme age, younger than 1 year and older than 70
99116	Anesthesia complicated by utilization of total body hypothermia
99135	Anesthesia complicated by utilization of controlled hypotension
99140	Anesthesia complicated by emergency conditions (specify)
99151	Moderate sedation services provided by the same physician or other qualified health care professional performing the diagnostic or therapeutic service that the sedation supports, requiring the presence of an independent trained observer to assist in the monitoring of the patient's level of consciousness and physiological status; initial 15 minutes of intraservice time, patient younger than 5 years of age
99152	Moderate sedation services provided by the same physician or other qualified health care professional performing the diagnostic or therapeutic service that the sedation supports, requiring the presence of an independent trained observer to assist in the monitoring of the patient's level of consciousness and physiological status; initial 15 minutes of intraservice time, patient age 5 years or older
99153	Moderate sedation services provided by the same physician or other qualified health care professional performing the diagnostic or therapeutic service that the sedation supports, requiring the presence of an independent trained observer to assist in the monitoring of the patient's level of consciousness and physiological status; each additional 15 minutes of intraservice time
99155	Moderate sedation services provided by a physician or other qualified health care professional other than the physician or other qualified health care professional performing the diagnostic or therapeutic service that the sedation supports; initial 15 minutes of intraservice time, patient younger than 5 years of age
99156	Moderate sedation services provided by a physician or other qualified health care professional other than the physician or other qualified health care professional performing the diagnostic or therapeutic service that the sedation supports; initial 15 minutes of intraservice time, patient age 5 years or older
99157	Moderate sedation services provided by a physician or other qualified health care professional other than the physician or other qualified health care professional

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue’s standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue’s Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Anesthesia Services and Moderate (“Conscious”) Sedation

performing the diagnostic or therapeutic service that the sedation supports; each additional 15 minutes of intraservice time

HCPCS

G0500

Moderate sedation services provided by the same physician or other qualified health care professional performing a gastrointestinal endoscopic service that sedation supports, requiring the presence of an independent trained observer to assist in the monitoring of the patient's level of consciousness and physiological status; initial 15 minutes of intra-service time; patient age 5 years or older

ICD-10 Diagnosis

All diagnoses

When services are Not Medically Necessary:

For the procedure codes listed above when criteria are not met.

Note: The following list of anesthesia service modifiers is for informational purposes:

CPT Physical Status Modifiers

- P1 A normal healthy patient (Class I)
- P2 A patient with mild systemic disease (Class II)
- P3 A patient with severe systemic disease (Class III)
- P4 A patient with severe systemic disease that is a constant threat to life (Class IV)
- P5 A moribund patient who is not expected to survive without the operation (Class V)

HCPCS Anesthesia Modifiers

- G8 Monitored anesthesia care (MAC) for deep complex, complicated, or markedly invasive surgical procedure
- G9 Monitored anesthesia care for patient who has history of severe cardio-pulmonary condition
- QS Monitored anesthesia care service

Discussion/General Information

Anesthesia services are provided by or under the supervision of a physician. Services consist of the administration of an anesthetic agent in various types of anesthesia.

Types of Anesthesia and Anesthesia Services

Anesthesia Service by the Surgeon: Anesthesia services personally furnished by the physician performing the surgical, therapeutic or diagnostic procedure are considered an integral component of the primary procedure. This may include local injections, regional blocks, and intravenous medication. General anesthesia administered and monitored by the surgeon is not considered medically appropriate.

Balanced Anesthesia: Anesthesia that uses a combination of drugs, each in an amount sufficient to produce its major or desired effect to the optimum degree and keep its undesirable or unnecessary effects to a minimum.

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Anesthesia Services and Moderate (“Conscious”) Sedation

Bier Block/Intravenous Regional Anesthesia (IVRA): Regional anesthesia produced by intravenous injection, used for surgical procedures on the arm below the elbow or the leg below the knee; performed in a bloodless field maintained by a pneumatic tourniquet that also prevents the anesthetic from entering the systemic circulation.

Brachial Plexus Block/Brachial Plexus Anesthesia: Regional anesthesia of the shoulder, arm, and hand by injection of a local anesthetic into the brachial plexus.

Caudal Block/Caudal Anesthesia: Regional anesthesia produced by injection of a local anesthetic into the caudal or sacral canal.

Epidural Block/Epidural Anesthesia: Regional anesthesia produced by injection of the anesthetic agent between the vertebral spines and beneath the ligamentum flavum into the epidural space.

General Anesthesia: A reversible state of unconsciousness and the inability to perceive pain, produced by anesthetic agents, with absence of pain sensation over the entire body and a greater or lesser degree of muscular relaxation; the drugs producing this state can be administered by inhalation, intravenously, intramuscularly, rectally, or via the gastrointestinal tract.

Inhalation Anesthesia: Anesthesia produced by the inhalation of vapors of a volatile liquid or gaseous anesthetic agent.

Intercostal Block/Intercostal Anesthesia: Anesthesia produced by blocking intercostal nerves with a local anesthetic.

Intranasal Anesthesia: Local anesthesia produced by insertion into the nasal fossae of pledgets soaked in a solution of an anesthetic agent which is effective after topical application, or by insufflation of a mixture of anesthetic gases or vapors through a tube introduced into the nose.

Intraoral Anesthesia: Anesthesia produced within the oral cavity by injection, spray, pressure, etc.

Intrathecal Anesthesia: Anesthesia produced by injection of an anesthetic solution into the subarachnoid space.

Intravenous Anesthesia/Intravenous Sedation (IV Sedation): Anesthesia produced by introduction of an anesthetic agent into a vein.

Local Anesthesia: Anesthesia confined to one area of the body.

Moderate (“Conscious”) Sedation: Involves the administration of medication with or without analgesia to achieve a state of depressed consciousness while maintaining the individual's ability to respond to stimulation. Moderate (“conscious”) sedation is administered by the surgeon or physician performing the procedure or an independent trained practitioner for the purpose of assisting the physician in monitoring the individual's level of consciousness and physiological status. It includes pre- and post-sedation evaluations, administration of the sedation and

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue’s standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue’s Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Anesthesia Services and Moderate (“Conscious”) Sedation

monitoring of the cardiorespiratory function. Cardiorespiratory functions monitored include heart rate, blood pressure and oxygen level.

Monitored Anesthesia Care (MAC): MAC was developed in response to the shift to providing more surgical and diagnostic services in an ambulatory, outpatient or office setting without the use of the traditional general anesthetic. Accompanying this, there has been a change in the provision of anesthesia services from the traditional general anesthetic to a combination of local, regional and certain consciousness altering drugs. This type of anesthesia is referred to as MAC if directly provided by anesthesia personnel. Based on the American Society of Anesthesiologists' (ASA) standards for monitoring, MAC should be provided by qualified anesthesia personnel (anesthesiologists or qualified anesthetists such as certified registered nurse anesthetists). These individuals must be continuously present to monitor and provide anesthesia care.

As described by the ASA's Position on Monitored Anesthesia Care (2018):

Monitored anesthesia care is a specific anesthesia service performed by a qualified anesthesia provider, for a diagnostic or therapeutic procedure. Indications for monitored anesthesia care include, but are not limited to, the nature of the procedure, the patient's clinical condition and/or the need for deeper levels of analgesia and sedation than can be provided by moderate sedation (including potential conversion to a general or regional anesthetic). Unlike monitored anesthesia care, moderate sedation is a proceduralist directed service which does not include a qualified anesthesia provider's periprocedural assessment and has the inherent limitations that are policy directed for the non-anesthesia qualified provider. Moderate sedation is a proceduralist directed service that may be governed by separate institutional policies.

Monitored anesthesia care includes all aspects of anesthesia care – a preprocedure assessment and optimization, intraprocedure care and postprocedure management that is inherently provided by a qualified anesthesia provider as part of the bundled specific service. During monitored anesthesia care, the anesthesiologist provides or medically directs a number of specific services, including but not limited to:

- Preprocedural assessment and management of patient comorbidity and periprocedural risk
- Diagnosis and treatment of clinical problems that occur during the procedure
- Support of vital functions inclusive of hemodynamic stability, airway management and appropriate management of the procedure induced pathologic changes as they affect the patient's coexisting morbidities
- Administration of sedatives, analgesics, hypnotics, anesthetic agents or other medications as necessary for patient safety
- Psychological support and physical comfort
- Provision of other medical services as needed to complete the procedure safely.

Monitored anesthesia care may include varying levels of sedation, awareness, analgesia and anxiolysis as necessary. The qualified anesthesiologist provider of monitored anesthesia care must be prepared to convert to general anesthesia and respond to the pathophysiology (airway and

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue's standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Anesthesia Services and Moderate (“Conscious”) Sedation

hemodynamic changes) of procedure and position in the management in induction of general anesthesia when necessary.

Regional Anesthesia: Anesthesia that involves the use of local anesthetic solutions(s) to produce circumscribed areas of loss of sensation. This includes spinal, epidural, nerve, field and extremity blocks. Spinal and epidural anesthesia is produced by injection of local anesthetic solution near the spinal canal, which interrupts sensation from the legs or abdomen.

Sacral Block/Sacral Anesthesia: Anesthesia produced by injection of a local anesthetic into the extradural space of the sacral canal.

Saddle Block Anesthesia: A type of sacral anesthesia produced in a region corresponding roughly with the area of the buttocks, perineum, and inner aspects of the thighs, by introducing the anesthetic agent low in the dural sac.

Spinal Anesthesia: Regional anesthesia produced by injection of a local anesthetic into the subarachnoid space around the spinal cord.

Standby Anesthesia: Anesthesia standby occurs when the anesthesiologist, or the CRNA, is available in the facility in the event he or she is needed for a procedure that requires anesthesia (e.g., available in the facility in case of obstetric complications - breech presentation, twins, and trial of instrumental delivery), but is not physically present or providing services. Standby anesthesia is not direct care (for instance, it is a standby service **without** direct hands-on contact).

Topical Anesthesia: Anesthesia produced by application of a local anesthetic directly to the area involved.

American Society of Anesthesiologists Levels of Sedation/Analgesia (ASA, 2019)

Minimal Sedation (Anxiolysis) is a drug-induced state during which patients respond normally to verbal commands. Although cognitive function and physical coordination may be impaired, airway reflexes, and ventilatory and cardiovascular functions are unaffected.

Moderate Sedation/Analgesia (“Conscious Sedation”) is a drug-induced depression of consciousness during which patients respond purposefully** to verbal commands, either alone or accompanied by light tactile stimulation. No interventions are required to maintain a patent airway, and spontaneous ventilation is adequate. Cardiovascular function is usually maintained.

Deep Sedation/Analgesia is a drug-induced depression of consciousness during which patients cannot be easily aroused but respond purposefully** following repeated or painful stimulation. The ability to independently maintain ventilatory function may be impaired. Patients may require assistance in maintaining a patent airway, and spontaneous ventilation may be inadequate. Cardiovascular function is usually maintained.

General Anesthesia is a drug-induced loss of consciousness during which patients are not arousable, even by painful stimulation. The ability to independently maintain ventilatory function

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue’s standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue’s Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Anesthesia Services and Moderate (“Conscious”) Sedation

is often impaired. Patients often require assistance in maintaining a patent airway, and positive pressure ventilation may be required because of depressed spontaneous ventilation or drug-induced depression of neuromuscular function. Cardiovascular function may be impaired.

Because sedation is a continuum, it is not always possible to predict how an individual patient will respond. Hence, practitioners intending to produce a given level of sedation should be able to rescue*** patients whose level of sedation becomes deeper than initially intended. Individuals administering Moderate Sedation/Analgesia (“Conscious Sedation”) should be able to rescue*** patients who enter a state of Deep Sedation/Analgesia, while those administering Deep Sedation/Analgesia should be able to rescue*** patients who enter a state of General Anesthesia.

*Monitored Anesthesia Care does not describe the continuum of depth of sedation, rather it describes “a specific anesthesia service performed by a qualified anesthesia provider, for a diagnostic or therapeutic procedure.” Indications for monitored anesthesia care include “the need for deeper levels of analgesia and sedation than can be provided by moderate sedation (including potential conversion to a general or regional anesthetic.”

**Reflex withdrawal from a painful stimulus is NOT considered a purposeful response.

***Rescue of a patient from a deeper level of sedation than intended is an intervention by a practitioner proficient in airway management and advanced life support. The qualified practitioner corrects adverse physiologic consequences of the deeper-than-intended level of sedation (such as hypoventilation, hypoxia and hypotension) and returns the patient to the originally intended level of sedation. It is not appropriate to continue the procedure at an unintended level of sedation.

References

Government Agency, Medical Society, and Other Authoritative Publications:

1. American Society of Anesthesiologists. ASA physical status classification system. Last amended October 23, 2019. For additional information visit the ASA website: <http://www.asahq.org>. Accessed on September 17, 2020.
2. American Society of Anesthesiologists. Continuum of depth of sedation: definition of general anesthesia and levels of sedation/analgesia. Last amended October 23, 2019. For additional information visit the ASA website: <http://www.asahq.org>. Accessed on September 17, 2020.
3. American Society of Anesthesiologists. Statement on granting privileges for administration of moderate sedation to practitioners who are not anesthesia professionals. Last amended October 26, 2016. For additional information visit the ASA website: <http://www.asahq.org>. Accessed on September 17, 2020.
4. American Society of Anesthesiologists. Statement on regional anesthesia. Last amended October 25, 2017. For additional information visit the ASA website: <http://www.asahq.org>. Accessed on September 17, 2020.
5. American Society of Anesthesiologists. Statement on granting privileges to non-anesthesiologist physicians for personally administering or supervising deep sedation. Last amended October 25, 2017. For additional information visit the ASA website: <http://www.asahq.org>. Accessed on September 17, 2020.

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue’s standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue’s Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Anesthesia Services and Moderate (“Conscious”) Sedation

6. American Society of Anesthesiologists. Practice guidelines for moderate procedural sedation and analgesia 2018: a report by the American Society of Anesthesiologists Task Force on Moderate Procedural Sedation and Analgesia, the American Association of Oral and Maxillofacial Surgeons, American College of Radiology, American Dental Association, American Society of Dentist Anesthesiologists, and Society of Interventional Radiology. March 2018. For additional information visit the ASA website: <http://www.asahq.org>. Accessed on September 17, 2020.
7. American Society of Anesthesiologists. Position on monitored anesthesia care. Last amended October 17, 2018. For additional information visit the ASA website: <http://www.asahq.org>. Accessed on September 17, 2020.

Index

Anesthesia Services
 Conscious Sedation
 General Anesthesia
 Moderate Sedation
 Monitored Anesthesia Care (MAC)
 Regional Anesthesia

History

Status	Date	Action
Reviewed	11/05/2020	Medical Policy & Technology Assessment Committee (MPTAC) review. Updated Description, Discussion/General Information and References sections. Reformatted Coding section.
Reviewed	11/07/2019	MPTAC review. Updated Discussion/General Information and References sections.
Revised	11/08/2018	MPTAC review. Removed statement on interventional pain management procedures from Clinical Indications section and moved to CG-MED-78 Anesthesia Services for Interventional Pain Management Procedures. Updated Discussion/General Information and References sections. Updated Coding section; removed CPT 01935, 01936, 01991, 01992.
Reviewed	09/13/2018	MPTAC review. Discussion/General Information and References sections updated.
Revised	11/02/2017	MPTAC review. Added a statement for when interventional pain management procedures are medically necessary. Updated Discussion and References sections. The document header wording updated from “Current Effective Date” to “Publish Date.” Updated Coding section with 01/01/2017 CPT changes; 01180, 01190, 01682 deleted 12/31/2017.
Reviewed	05/04/2017	MPTAC review. Formatting updated in Clinical Indications section. Description and References sections updated.
	01/01/2017	Updated Coding section with 01/01/2017 CPT and HCPCS changes; removed codes 99143, 99144, 99145, 99148, 99149, 99150 deleted 12/31/2016 and codes for nerve blocks which are not used for anesthesia during procedures.

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue’s standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue’s Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Anesthesia Services and Moderate (“Conscious”) Sedation

Reviewed	05/05/2016	MPTAC review. References section updated.
	01/01/2016	Updated Coding section with 01/01/2016 CPT changes, removed 64412 deleted 12/31/2015; also removed ICD-9 codes.
Reviewed	05/07/2015	MPTAC review. Description, Discussion and References sections updated.
	01/01/2015	Updated Coding section with 01/01/2015 CPT changes; removed 00452, 00622, 00634 deleted 12/31/2014.
Reviewed	05/15/2014	MPTAC review. References section updated.
Reviewed	05/09/2013	MPTAC review. References updated.
Reviewed	05/10/2012	MPTAC review. References updated.
Reviewed	05/19/2011	MPTAC review. References updated.
Reviewed	05/13/2010	MPTAC review. Discussion and References updated.
	01/01/2010	Updated Coding section with 01/01/2010 CPT changes; removed CPT 01632 deleted 12/31/2009.
Reviewed	05/21/2009	MPTAC review. Discussion, Coding and References updated. Place of service section removed.
Revised	05/15/2008	MPTAC review. Added a statement for when anesthesia services are not medically necessary. References and Appendix updated. Coding updated with 01/01/2008 CPT updates; removed CPT 01905 deleted 12/31/2007.
Reviewed	05/17/2007	MPTAC review. References updated.
Revised	06/08/2006	MPTAC review. Document title revised. Term conscious sedation updated to moderate sedation per ASA guidelines. Updated definition of MAC per ASA guidelines. Indications for anesthesia services during gastrointestinal endoscopic procedures removed. References updated.
Revised	03/23/2006	MPTAC review. Updated language for regional anesthesia. Revision per recommendation from American Society of Anesthesiologists.
	01/01/2006	Updated coding section with 01/01/2006 CPT/HCPCS changes.
Revised	09/22/2005	MPTAC review. Revision based Pre-merger Anthem and Pre-merger WellPoint Harmonization.

Pre-Merger Organizations	Last Review Date	Document Number	Title
Anthem, Inc. WellPoint Health Networks, Inc.	04/28/2005	Definition vii	No document Anesthesia Services

Appendix

American Society of Anesthesiology Physical Status Classifications:

ASA I A normal healthy patient

ASA II A patient with mild systemic disease

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue’s standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue’s Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

Anesthesia Services and Moderate (“Conscious”) Sedation

ASA III A patient with severe systemic disease

ASA IV A patient with severe systemic disease that is a constant threat to life

ASA V A moribund patient who is not expected to survive without the operation

ASA VI A declared brain-dead patient whose organs are being removed for donor purposes

This Clinical UM Guideline is intended to provide assistance in interpreting Healthy Blue’s standard Medicaid benefit plan. When evaluating insurance coverage for the provision of medical care, federal, state and/or contractual requirements must be referenced, since these may limit or differ from the standard benefit plan. In the event of a conflict, the federal, state and/or contractual requirements for the applicable benefit plan coverage will govern. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Clinical UM Guideline is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue’s Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.