

<b>Subject:</b>	MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications	<b>Publish Date:</b>	09/27/2023
<b>Document #:</b>	MED.00057	<b>Last Review Date:</b>	08/10/2023
<b>Status:</b>	Reviewed		

## Description/Scope

This document addresses magnetic resonance imaging (MRI) guided high intensity focused ultrasound (HIFU) ablation, also known as magnetic resonance guided focused ultrasound (MRgFUS), when used to treat any non-oncologic indications, including but not limited to uterine fibroids, essential tremor (ET), or benign prostatic hyperplasia (BPH). The ultrasound beam penetrates through the soft tissues and can be focused to targeted sites, using MR for guidance and monitoring.

**Note:** Please see the following related documents for additional information:

- CG-MED-81 Ultrasound Ablation for Oncologic Indications
- CG-SURG-28 Transcatheter Uterine Artery Embolization
- CG-SURG-91 Minimally Invasive Ablative Procedures for Epilepsy
- SURG.00026 Deep Brain, Cortical, and Cerebellar Stimulation

## Position Statement

### Medically Necessary

Unilateral focused ultrasound thalamotomy is considered **medically necessary** for the treatment of adults with essential tremors when **all** of the following criteria are met:

1. Moderate to severe tremor of the hand (as defined by a score of 2 or greater on the clinical rating scale for tremor [CRST]); **and**
2. Failure of 2 or more tremor suppressant medications, as evidenced by persistent moderate to severe tremors, intolerable side effects of drug therapy or contraindications; **and**
3. At least 1 of the medications failed was a first line agent (propranolol or primidone).

### Investigational and Not Medically Necessary:

MRI guided high intensity focused ultrasound ablation is considered **investigational and not medically necessary** when the criteria above are not met and for all other non-oncologic indications.

## Rationale

### *Essential Tremor (ET)*

The first line treatment for ET is pharmacotherapy, with propranolol and primidone considered as first line medication (Elias, 2016). Neurosurgical intervention targeted at the nucleus ventralis intermedius of the thalamus

This Medical Policy provides assistance in understanding Healthy Blue's standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications**

---

can be considered for individuals who have failed medication therapy. The targeted tissue connects the cerebellum with cortical motor pathways. Radiofrequency thalamotomy and deep brain stimulation have been used to disrupt this pathway and suppress tremors. Acceptance of these procedures by those with ET has generally been low based upon the reluctance to undergo the invasive procedures.

In July 2016, the U.S. Food and Drug Administration (FDA) approved the ExAblate Neuro<sup>®</sup> system (InSightec, Inc., Dallas, TX) as a treatment of idiopathic ET in adults aged 22 or older whose tremor has failed pharmacological treatment. ExAblate Neuro uses focused ultrasound to induce a unilateral thalamic lesion (thalamotomy) when the ventralis intermedius has been identified and is accessible for ablation by the device. The intent of treatment is to reduce an individual's ET and increase motor function.

Elias and colleagues (2013) reported on the results of a feasibility trial for the ExAblate Neuro unilateral transcranial MRI-guided focused ultrasound thalamotomy to treat medication-refractory ET. The study defined medication refractory tremor as failure of treatment with at least two trials of full-dose therapeutic medication, one of which had to be a first line therapy (propranolol or primidone). In this uncontrolled pilot study, 15 individuals with severe medication-refractory ET underwent a single treatment session using the ExAblate Neuro system. Assessments were performed at baseline, 1 day, 1 week, 1 month, 3 months and 12 months following treatment, with the change in hand tremor score at 3 months being the primary clinical outcome. Hand tremor was scored using a summation of eight items which graded hand tremor and ability to perform tasks. Higher scores denoted more severe tremor, and the maximum score was 32. There was significant improvement in contralateral hand tremor from baseline ( $20.4 \pm 5.2$ ) to 3 months ( $4.3 \pm 3.5$ ) and 12 months ( $5.2 \pm 4.8$ ). No significant difference in ipsilateral hand tremor scores from baseline to 12 months was found. Paresthesias of the face or fingers were the most common reported side effects, with 4 individuals reporting persistent paresthesias. The authors recommended that larger studies were needed to evaluate potential associated cognitive impairment and include a comparison to the current standard therapy.

Subsequent to the above pilot study, Elias and colleagues (2016) published the results of a prospective, sham-controlled, double-blind, randomized trial of 76 participants, which evaluated MRI-guided focused ultrasound thalamotomy treatment of moderate to severe medication-refractory ET. Medication refractory treatment was defined as tremor refractory to at least 2 agents, with at least one being propranolol or primidone. Participants were assigned in a 3:1 ratio to undergo either the active treatment, MRI-guided focused ultrasound thalamotomy, or sham treatment. Following evaluation of the primary endpoint at 3 months, the individuals in the sham group could cross over to the active treatment group. The change in hand tremor scores from baseline to 3 months was defined as the primary efficacy outcome measure. The hand tremor score was based upon components of the Clinical Rating Scale for Tremor (CRST) related to hand tremor, with higher scores indicating more severe tremor. At 3 months, the mean score in the contralateral hand in the active treatment group improved by 47% compared to the sham group mean score improvement of 0.1% ( $18.1 \pm 4.8$  to  $9.6 \pm 5.1$  versus  $16.0 \pm 4.4$  to  $15.8 \pm 4.9$ ; between group change 8.3 points; 95% confidence interval [CI], 5.9-10.7;  $p=0.001$ ). At 12 months, the significant improvement over baseline persisted. There was no significant change in the tremor score in the ipsilateral hand compared to baseline. The results were similar in the sham crossover group; 19 participants who crossed over to active treatment reported a significant improvement in contralateral hand tremor at 3 and 6 months respectively ( $16.5 \pm 4.2$  to  $7.4 \pm 3.9$  and  $16.5 \pm 4.2$  to  $8.0 \pm 3.9$ ). A total of 74 neurological adverse events occurred in 56 individuals who underwent active treatment, including 38% with sensory alteration and 36% with cerebellar deficits such as dysmetria and ataxia and other gait disturbance, which persisted to 12 months at 14% and 9%, respectively.

---

This Medical Policy provides assistance in understanding Healthy Blue's standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

---

**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications**

---

Several follow-up studies have evaluated durability of MRI-guided focused ultrasound thalamotomy 2 to 3 years following treatment. Chang and colleagues (2018) reported on the 2-year follow-up outcomes of the Elias 2016 randomized controlled trial (RCT). The mean hand tremor score initially improved by 55% (from  $19.8 \pm 4.9$  at baseline to  $8.6 \pm 4.5$ ) at 6 months post procedure. At 2 years, the mean hand tremor motor score was improved by 56% over baseline ( $8.8 \pm 5.0$ ; change in the score from baseline to 2 years, 11 points). At 3 years post-procedure, Halpern and colleagues (2019) evaluated outcomes of this same group. The median hand tremor motor score was stable at 56% improvement over baseline (median score of 8). In a retrospective review, Meng and associates (2018) assessed the 2-year outcomes of 37 individuals who underwent unilateral MRgFUS thalamotomy to treat moderate to severe medically refractory ET. A 42.4% (95% CI, 32.0%-52.9%) improvement in the baseline dominant tremor score ( $20.3 \pm 5.0$ ) was maintained at 2 years (43.4%: 95% CI: 27.8%-59.0%). At 1 year post-treatment, 45.7% of the individuals had significant tremor improvement; this had decreased to 35.3% at 2 years (Halpern, 2019). These follow-up studies report substantial drop-out rates, introducing the potential for retention bias into the results. Additional studies with follow-up from 3 to 5 years post-procedure report showed sustained improvement from baseline (Cosgrove, 2022; Halpern, 2019; Park, 2019; Sinai, 2019).

The International Parkinson and Movement Disorder Society published an evidence-based review of ET treatments (Ferreira 2019). The task force noted that unilateral MRgFUS thalamotomy is “likely efficacious” (evidence suggests, but is not sufficient to show, that the intervention has a positive effect on studied outcomes). The task force concluded that unilateral MRgFUS thalamotomy is “possibly useful” for clinical practice. The society’s recommendation is based on the Elias (2016) RCT.

Miller and colleagues (2022) reported on the prevalence of worsening tremor over time following MRgFUS treatment of ET. This is the phenomenon noted above in Louis’s analysis of the pivotal trial published by Elliot et al. Miller’s meta-analysis included 17 prospective studies, 3 retrospective studies and 1 RCT. Tremor was evaluated using hand tremor scores (HTS), CRST scores, or Quality of Life in Essential Tremor Questionnaire (QUEST) using pooled reported effects. The analysis showed ongoing treatment benefit but decreasing treatment effect from 3 to 12 months and 24 months following treatment. The authors noted that this diminishing effect might be due to heterogeneity within the studies, disease progression, or a true waning effect over time. Diminished effects have also been reported with DBS, either due to disease progression or habituation. At this time, there are no studies which directly compare MRgFUS and DBS.

The American Society for Stereotactic and Functional Neurosurgery (ASSFN) position statement on MRgFUS (Pouratian, 2020) notes the following indications when using MRgFUS as a treatment option of ET:

1. Confirmed diagnosis of ET.
2. Failure to respond to, intolerance of, or medical contraindication to use of at least 2 medications for ET, 1 of which must be a first-line medication.
3. Appendicular tremor that interferes with quality of life (QoL) based on clinical history.
4. Unilateral treatment.

The ASSFN also notes the following contraindications to MRgFUS therapy:

1. Bilateral MRgFUS thalamotomy.
2. Contralateral to a previous thalamotomy.
3. Cannot undergo magnetic resonance imaging (MRI) because of medical reasons.
4. Skull density ratio (ratio of cortical to cancellous bone) is  $<0.40$

---

This Medical Policy provides assistance in understanding Healthy Blue’s standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue’s Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications**

---

The ASSFN statement also notes that there is insufficient evidence to support treatment of tremors of the head, voice or neck with MRgFUS.

In 2020, Giordano and associates published a systematic review comparing unilateral MRgFUS thalamotomy to unilateral and bilateral DBS. Studies reporting on the treatment of drug-refractory ET using DBS (n=37) or MRgFUS (n=7) were included. There were no prospective randomized studies directly comparing MRgFUS to DBS. A total of 1202 individuals were included in this study's DBS group and 477 individuals were included in the MRgFUS group. At 14.4-16.6 months follow-up, the improvement in quality of life was significantly greater in the MRgFUS group (61.9%) compared to the DBS group (52.5%). During that follow-up period, there was a significantly higher improvement in tremor severity in the DBS group (60.1%) compared to the MRgFUS group (55.6%). While a subgroup analysis showed no differences in tremor severity between unilateral DBS therapy and MRgFUS therapy, bilateral DBS therapy was superior to both unilateral treatments. The authors asserted that bilateral DBS is the current gold standard treatment for medication-resistant ET. Bilateral staged MRgFUS thalamotomy is currently undergoing feasibility testing. MRgFUS and DBS have different complication patterns. MRgFUS is associated with a higher prevalence of gait disturbances/muscle problems, nausea and paresthesias. DBS is associated with a higher prevalence of speech disturbances and local adverse symptoms. Persistent complications are more frequent with MRgFUS than with DBS therapy. This may be due to permanent tissue destruction induced by MRgFUS compared to DBS which can be reprogrammed or turned off.

*MRgFUS for ET Summary*

Radiofrequency or stereotactic radiosurgery thalamotomy (Gamma Knife) or DBS have historically been the standard treatments for medically refractory ET. These procedures are associated with risks including intracerebral hemorrhage, infections, and hardware malfunction. Adoption of these modalities by individuals with medically refractory ET has been limited due to the invasiveness of these procedures and their perceived risks. MRgFUS has become widely accepted by the medical community as an effective minimally invasive therapy for individuals who are not candidates for, or refuse to undergo, a more invasive procedure.

*Parkinson's Disease and other non-ET movement disorders*

In an initial pilot investigation, Bond and associates (2017) evaluated the safety and efficacy of focused ultrasound thalamotomy for the treatment of medically refractory, tremor-dominant Parkinson disease (TDPD). Like ET, medically refractory TDPD can be treated with DBS or thalamic lesioning. In a double-blind, sham-controlled, pilot RCT, 27 individuals were treated with focused ultrasound (FUS) thalamotomy (n=20) or sham procedure (n=7). After 3 months, individuals in the sham procedure group were offered open-label treatment and 6 individuals underwent FUS thalamotomy. The primary outcome was the change from baseline to 3 months in the treated upper limb tremor subscore. Hand tremor improved 62% (interquartile range [IQR] 22%-79%) from a baseline of 17 points (IQR 10.5-27.5) following FUS thalamotomy and 22% (IQR -11% to 29%) from a baseline of 23 points (IQR 14-27) after sham procedures (p=0.04). Adverse events occurred in approximately 35% of all individuals treated and included finger paresthesia, ataxia, and orofacial paresthesia. At 1 year, in those individuals in the initial treatment group that were available for evaluation (14/30), 13 individuals reported a positive outcome. While this initial pilot study reported positive results, this was a small study which did not reach its planned enrollment size and lost a significant portion of the treatment group at 1 year follow-up. In addition,

---

This Medical Policy provides assistance in understanding Healthy Blue's standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications**

---

the authors noted a potential confounder in allowing medication dose changes during the trial. Larger trials comparing FUS thalamotomy to standard treatment with longer follow-up are needed.

Altinel and colleagues (2019) conducted a systematic review and meta-analysis comparing lesion surgery to DBS to treat tremor related to Parkinson's Disease, ET or multiple sclerosis. While the analysis included 15 randomized studies, only 2 studies used MRgFUS to create the lesion. The duration of follow-up in both studies was limited to 3 months. A separate analysis found no difference in tremor severity improvement between MRgFUS, DBS and other types of lesion surgery. Over the short-term 3-month follow-up period, MRgFUS was associated with higher QOL scores than DBS. This analysis was limited by heterogeneity (tremor etiology, type of lesion surgery), limited follow-up times and lack of direct comparison groups.

In 2018, Schreglmann and associates evaluated efficacy and the prevalence of persistent side effects of different lesioning techniques in the treatment of ET, Parkinson's disease, dystonic tremor, multiple sclerosis or lesions to the midbrain or cerebellar structures. ET was treated with MRgFUS, Gamma Knife or radiofrequency in 6 retrospective and 7 prospective studies. The primary outcome was the change in upper limb tremor severity from baseline to follow-up, with the selected follow-up time-point as the point with the largest number of individuals retained. As the follow-up times varied, the authors controlled for an effect on follow-up duration on the effect size. The authors reported that the duration of follow-up did not have a significant influence on treatment effect size. There were no significant differences over the studied time periods with regard to the mean effect on tremor severity or the rate of persistent side effects. The authors concluded that head-to-head comparisons between DBS and MRgFUS are needed to further evaluate tremor treatment and noted:

Nevertheless, this systematic review also shows how limited the evidence base is in particular for MRIgFUS ablation in conditions other than ET. It therefore highlights the need for adequately designed prospective trials to support the existing data on safety and efficacy for established targets such as V.im. and of recently rediscovered targets within the PSA. Before that, the indiscriminate application of incisionless interventions to novel indications could potentially harm the further development of this fascinating technique.

Krishna and associates (2023) evaluated the safety and efficacy of focused ultrasound ablation (FUSA) of the globus pallidus internus in individuals with Parkinson's disease. Individuals with Parkinson's disease and dyskinesias or motor fluctuations and motor impairment in the off-medication state were eligible to be included in this multicenter, prospective, double-blind, randomized, sham-controlled trial. Trial eligibility was determined by scores on the Movement Disorders Society–Unified Parkinson's Disease Rating Scale, part III (MDS-UPDRS III), for the treated side in the off-medication state or in the score on the Unified Dyskinesia Rating Scale (UDysRS) in the on-medication state. Participants were randomized to receive FUSA of the globus pallidus internus (n=69) or a sham procedure (n=25). The primary outcome was a response- a decrease or improvement of at least 3 points on either of the cited scales on the treated side without a clinically meaningful worsening in either scale at 3 months. A total of 65/69 (94%) participants in the treatment group and 22/25 (88%) in the sham group completed the 3-month assessment. At 3 months, approximately one-third of participants in the FUSA group were classified as having a response resulting in improved motor function and reduced dyskinesia, one-third were classified as non-responders. Approximately one-third of participants in the sham group had a response. After the 3-month evaluation, 20 individuals in the sham group elected to undergo active treatment. The 3-month outcomes of the cross-over group were similar to the initial active treatment group. The authors of the study concluded:

---

This Medical Policy provides assistance in understanding Healthy Blue's standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications**

---

The procedure has recently been approved by the FDA for unilateral treatment only, whereas most patients with Parkinson's disease have bilateral motor signs and symptoms. There are limitations of the procedure in a condition with bilateral manifestations. Unlike deep-brain stimulation, there is no opportunity to treat progressive worsening of motor symptoms. Owing to the lack of ability to modify the lesion after the procedure, operators must prioritize safety during treatment. With deep-brain stimulation, adjustments in programming can attempt to improve the balance between a reduction in motor symptoms and a reduction in stimulation-related side effects, even years after implantation.

Schrag (2023) commented that the results of the Krishna (2023) study were promising, but noted:

given the nonreversible nature of the intervention and the progressive nature of the disease, it will be important to establish whether improvements in motor complications are maintained over longer periods and whether treatment results in improved overall functioning and quality of life for patients.

***BPH***

HIFU ablation is a minimally invasive procedure using a transrectal ultrasound probe to image the prostate and deliver timed bursts of heat to create coagulation necrosis in a targeted area without harming adjacent healthy tissue (Leslie, 2006). Schatzl and colleagues (2000) compared the efficacy of transurethral resection of the prostate (TURP) to four less invasive treatment options including HIFU in a small clinical trial. Randomization was attempted but could not be carried out because treatment options for each participant were limited based on specific characteristics such as prostate size, prostatic calcifications and middle lobes. The individuals who received HIFU tended to have smaller prostates and less severe symptoms than those who received TURP. A second study reported by Madersbacher and colleagues (2000) attempted to determine the long-term outcome after HIFU therapy for individuals with lower urinary tract symptoms (LUTS) due to BPH. The data collected between June 1992 and March 1995 indicated that HIFU therapy for BPH, at least in its present form, did not "stand the test of time," as 43.8% of individuals had to undergo TURP within 4 years after initial therapy. Additional long-term studies are warranted to reliably assess the role of HIFU as an established alternative to standard treatments for BPH. In recent years, few trials evaluating the use of HIFU in BPH have been published.

***Uterine Fibroids***

Stewart and colleagues (2003) studied the safety and feasibility of MR-guided HIFU (referred to as focused ultrasound surgery [FUS] in this study) in a case series of 55 women with symptomatic uterine fibroids who underwent HIFU treatment and, in some cases, a planned hysterectomy within 1 month after the ultrasound treatment (n=28). In the latter group, hysterectomy specimens provided pathologic evidence of accurate levels of thermal energy delivered and revealed 3 times the volume of necrotic fibroid tissue than was targeted for treatment. Reported side effects related to HIFU were minimal. Authors concluded that "...correlation of both treatment and patient parameters with the surrogate endpoint of post-FUS MRI and, ultimately, symptom reduction will be necessary in future studies to optimize therapy for individual women."

In the study designed for FDA approval of the ExAblate® 2000 (InSightec, Ltd, Dallas, TX), Stewart and colleagues studied 109 women treated with MR-guided HIFU and 83 women treated with abdominal

---

This Medical Policy provides assistance in understanding Healthy Blue's standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications**

---

hysterectomy. The initial article published in 2004 and the follow-up article published in 2006, only reported the outcomes for the group treated with HIFU. The study's primary outcome was change in the symptom severity score (SSS) that is part of the validated Uterine Fibroid Symptom Quality of Life Questionnaire (UFS-QOL). Symptom severity was measured on a 0 (less severe) to 100 (most severe) scale with eight questions relevant to bulk and bleeding symptoms. At the 6-month follow-up, 71% of the HIFU group achieved a 10-point or greater reduction in SSS which decreased to 51% at 12 months. It was unclear what value represented a clinically meaningful change in SSS. Furthermore, 21% of those treated by HIFU needed additional surgical treatment, and 4% underwent a repeat MRI guided HIFU within 12 months of the first treatment.

Stewart and colleagues (2007) published results from three phase III trials and one post-marketing study. A total of 416 women were enrolled who received HIFU. Quality of life outcomes, measured again by the SSS of the UFS-QOL, were assessed for 359 women who were available at the 24-month follow-up. Clinical endpoints of the trials included uterine shrinkage, the need for additional fibroid treatment and the time to additional fibroid treatment. The study found a relationship between the non-perfused volume ratio and the probability of undergoing additional fibroid treatment. There was significantly greater improvement in women who had a more complete ablation; however, for women with minimal ablation, the need for additional treatment/procedures was high.

Fennessy and colleagues (2007) evaluated MR-guided HIFU using different treatment protocols. Results from nonrandomized, consecutive participants treated with the original protocol (33% of fibroid volume with a maximum treatment time of 120 minutes, n=96) were compared to a modified protocol (50% treatment volume, 180-minute maximum treatment time, and a second treatment if within a 14-day period, n=64). In the original protocol group, the non-perfused (effectively treated) area was calculated at 17% of fibroid volume compared with 26% of fibroid volume with the modified protocol group. Overall, symptom severity was reported to have decreased from a score of 62 at baseline to 33 at 12 months, with fewer participants in the modified group choosing alternative treatment (28% versus 37%). Out of the 160 participants initially treated, 55 from the original treatment protocol and 21 from the modified protocol group were evaluated at the 12-month follow-up.

In 2009, Taran and colleagues reported outcomes for the hysterectomy group in the 2006 Stewart study. The Taran article did not use the original primary outcome measure (SSS scores) and instead reported findings using the SF-36 quality of life measure as well as safety data. A significantly higher proportion of women in the hysterectomy group (82 of 83, 99%) reported at least one adverse event compared to women in the MR-guided HIFU group (88 of 109, 81%). Pain or discomfort, adverse events associated with the gastrointestinal tract, dermatological system, nervous system, and cardiovascular system were significantly more common in the hysterectomy group. However, a similar proportion reported a serious adverse event, 9 of 109 (8%) in the HIFU group and 8 of 83 (10%) in the hysterectomy group. The authors concluded that the results of their study showed that HIFU treatment of uterine fibroids leads to clinical improvement with fewer significant clinical complications and adverse events compared to hysterectomy during the 6-month follow-up period. The authors did acknowledge important limitations of the study--factors that were consistent with more severe disease. The women undergoing hysterectomy had increased body mass index (BMI), were less likely to be Caucasian, had higher symptom severity scores and had an increased use of medication for fibroid-related symptoms. They also pointed out the possibility of potential data collection bias between the novel-treatment group and the control group. Although this study proposes HIFU as a treatment alternative to hysterectomy for symptomatic fibroids, it did not demonstrate clinical efficacy when compared to the current accepted treatments of uterine fibroids.

---

This Medical Policy provides assistance in understanding Healthy Blue's standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications**

---

Kim and colleagues (2011) reported a 3-year follow-up on a prospective study of 40 women with symptomatic fibroids. A total of 51 fibroids were treated with MR-guided HIFU. Clinical assessments were obtained at 3 months, 6 months, and 1, 2, and 3 years after HIFU treatment, as well as the SSS from the UFS-QOL. An MRI was performed at each follow-up to assess the efficacy of the treatment at 6 months, 1 year, 2 years, and 3 years. The mean baseline volume of treated fibroids was 336.9 cm<sup>3</sup>. The mean improvement scores for transformed SSS was 47.8 (p<0.001) and for transformed UFS-QOL was 39.8 (p<0.001) at 3 years. The mean volume decrease in treated fibroids was 32.0% (p<0.001) and in the uterus, the volume decrease was 27.7% (p<0.001) at 3 years. There were no complications. The authors noted that although these results are preliminary, MR-guided HIFU for the treatment of uterine fibroids may be an acceptable treatment option.

Ikink (2013) conducted a nonrandomized uncontrolled study to assess the clinical efficacy of MR-guided ultrasound techniques for the treatment of symptomatic uterine fibroids (n=46 premenopausal women). Clinical outcomes were measured using the SSS of the UFS-QOL questionnaire. Results showed that HIFU treatment resulted in a significant reduction in fibroid volume compared with baseline measurements (p<0.001), corresponding to a volume reduction of about 29%. In addition, 25 of 46 women (54%) reported a significant improvement in mean transformed SSS compared with baseline values (p<0.001). This study was also characterized by significant limitations, including lack of randomization and lack of a control arm.

Clark and colleagues (2014) conducted a systematic review of the efficacy of MRgFUS, specifically on its performance preserving fertility in women. A total of 10 studies, representing 589 women, were chosen for inclusion in the meta-analysis. Study inclusion criteria included a report of mean SSS at baseline and 6-month follow-up. The overall mean improvement in SSS 6 months after MRgFUS was estimated at 31.0% (95% CI, 23.9-38.2%). Authors concluded that, "Given the minimally invasive approach, MRgFUS could become the treatment of choice for patients desiring future fertility; however, further investigation is needed."

Another 2014 systematic review completed by Gizzo and colleagues evaluated the safety, feasibility, indications, complications, impact on UFS-QOL and fertility associated with myomectomy performed by MRgFUS. This review included 38 studies and a total of approximately 2500 women. While the authors concluded that MRgFUS was a safe and efficient technique, it was noted that comparisons with other traditional techniques have not yet been performed.

Pron (2015) reviewed the published evidence related to MRgHIFU to treat uterine fibroids. The evidence used in the review included two systematic reviews, two RCTs, 45 cohort study reports, and 19 case reports. This evaluation did not include any randomized trials comparing MRgFUS to other guidance methods, other minimally invasive treatments, or surgeries for symptomatic uterine fibroids. The author concluded MRgHIFU can be a safe and effective non-invasive, uterine preserving treatment alternative to hysterectomy for individuals failing medical therapy. However, the author also noted there is limited information regarding treatment durability. In addition, there is a lack of existing evidence comparing MRgHIFU to the current standard treatment.

Well-designed clinical studies evaluating the safety and efficacy of MRgFUS using relevant outcome measures are lacking. Studies have generally lacked comparison to the current accepted treatments of uterine fibroids. Additionally, based on the prevalence of uterine fibroids, a relatively small number of women have been studied in clinical trials. The published evidence regarding MRgFUS does not adequately address the potential for regrowth of treated uterine fibroids over time, particularly beyond 3 to 5 years. In order to demonstrate MRgFUS

---

This Medical Policy provides assistance in understanding Healthy Blue's standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.



**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications**

as a safe and effective treatment option for uterine fibroids, well-designed RCTs with sufficient follow-up periods and appropriate clinical outcome measures are needed to compare therapy with alternative treatments.

The American College of Radiology (ACR) Appropriateness Criteria® Treatment of Uterine Leiomyomas (2017) states that:

To date, there is little long-term information on the efficacy of this technology. It has been reported that myomas treated with HIFU have nearly 50% volume reduction at 1 year, but viable cells are present at biopsy in nearly 26% of specimens. Funaki et al. report a 24-month volume reduction of 40% with significant symptomatic improvement at 6 months that remained stable at 24-month follow-up. In a multicenter trial, Stewart et al. demonstrated significant reduction in fibroid-related symptoms in 70% of patients at 6 months and 51% of patients at 12 months.

The American College of Obstetricians and Gynecologists (ACOG) stated in their practice bulletin, Management of Symptomatic Uterine Leiomyomas (2021):

Limited, low-quality data suggest that magnetic resonance-guided focused ultrasound and high-intensity focused ultrasound are associated with a reduction in leiomyoma and uterine size. However, small randomized comparative trial data suggest that compared with UAE, magnetic resonance-guided focused ultrasound is associated with less improvement in symptoms and quality-of-life measures and a higher risk of reintervention.

**Other Conditions**

MRgFUS is being studied for a variety of conditions, including but not limited to benign thyroid nodules, chronic neuropathic pain, desmoid tumors, obsessive compulsive disorder or primary hyperparathyroidism (Chung, 2020; Jeanmonod, 2012; Jung, 2015; Kovatcheva, 2014; Martin, 2009; Monteith, 2016). These preliminary studies are in the early stages of evaluating the feasibility, efficacy and safety of the use of MRgFUS for the treatment of these conditions.

**Background/Overview****BPH**

BPH, the nonmalignant growth of the smooth muscle and epithelial cells within the prostate, is a common age-related manifestation. While almost all men will develop histologic BPH, this condition will not require treatment unless it is associated with subjective symptoms, such as LUTS. LUTS symptoms can be divided into two categories. Obstructive symptoms include hesitancy, straining, weak flow, prolonged voiding, partial or complete urinary retention or overflow incontinence. Irritative symptoms include frequency, urgency with urge incontinence, nocturia, and painful urination or voiding small amount (Roehrborn, 2005). The primary goal of treatment is to alter disease progression and prevent complications (AUA, 2014). If BPH has advanced to the point of causing obstruction, interventions may be aimed at removing excessive tissue and relieving the obstruction. HIFU has been proposed as a means of removing excessive prostate tissue.

This Medical Policy provides assistance in understanding Healthy Blue's standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

## Medical Policy

### MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications

#### ET

ET is a progressive, disabling movement disorder, affecting as much as 4% of the population (Elias, 2013). ET does not reduce life expectancy but does adversely affect QOL. While there does appear to be a familial component, the cause of this disorder is not known. Medication is the initial treatment when the symptoms interfere with function and QOL, with propranolol and primidone considered first-line agents. Second-line agents, such as gabapentin or carbamazepine are considered less effective. Approximately 30- 50% of individuals treated for ET cannot tolerate or fail medication therapy. Surgical treatments include radiofrequency thalamotomy, surgical resection and deep brain stimulation. MRgFUS is considered a minimally invasive treatment option. Similar to radiofrequency thalamotomy and surgical resection, MRgFUS creates a thalamic lesion which can reduce tremor, but can also result in permanent neurologic deficits (Elias, 2016). While creation of a lesion does reduce tremor and larger lesions can result in more enduring efficacy, larger lesions have a higher incidence of side effects (Elias, 2016).

Tremor severity is quantified using a tremor rating assessment scale. Fahn–Tolosa–Marin Clinical Rating Scale and the Essential Tremor Rating Assessment Scale (TETRAS) rate tremors on a scale from 0 to 4. The initial tool, the Fahn–Tolosa–Marin Tremor clinical rating scale was not specific to ET. TETRAS, developed specifically for ET, correlates strongly with the Fahn–Tolosa–Marin Clinical Rating Scale. TETRAS was developed for use to evaluate ET severity in clinical trials and to monitor disease progression. The following table excerpt represents the upper limb tremor as defined by the Essential Tremor Rating Assessment Scale (Elble, 2012):

Rating	Upper limb tremor
0	No tremor
1	Barely visible
1.5	< 1 cm
2	1 - < 3 cm
2.5	3 - < 5 cm
3	5 - < 10 cm
3.5	10 - < 20 cm
4	≥ 20 cm

#### Uterine Fibroids

The cause of fibroid tumors, also known as leiomyomas, of the uterus is unknown. However, it is suggested that fibroids may enlarge with estrogen therapy (such as oral contraceptives) or with pregnancy. Fibroid growth seems to depend on regular estrogen stimulation, and rarely affects women younger than 20 years of age or postmenopausal women. As long as a woman with fibroids is menstruating, the fibroids will probably continue to grow, although growth is usually quite slow.

Hysterectomy and various myomectomy procedures are considered the gold standard of treatment. However, there has been a longstanding research interest in developing minimally invasive alternatives. There has been interest in using HIFU treatment as a noninvasive approach to ablation of uterine fibroids. Treatment involves the use of focused high-intensity convergent ultrasound beam which increases the temperature within the targeted area to 60-95°C to destroy tissue without causing damage to adjacent tissue. During the procedure, individuals are typically

---

This Medical Policy provides assistance in understanding Healthy Blue's standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications**

---

placed under conscious sedation with or without epidural anesthesia, although general anesthesia may be used. Proposed advantages of HIFU include the noninvasive nature of the procedure that spares surrounding tissue, reducing postoperative morbidity, and hastening recovery.

The U.S. Food and Drug Administration (FDA) approved, via the Premarket Application (PMA) process, the ExAblate 2000 System for ablation of uterine fibroid tissue in pre- or peri-menopausal women with symptomatic uterine fibroids who desire a uterine sparing procedure. The 2004 FDA approval letter stated that women must have a uterine gestational size of less than 24 weeks and must have completed childbearing (FDA, 2004).

**Definitions**

**Benign prostate hyperplasia (BPH):** A condition that causes an increase in the size of the prostate gland in men, commonly causing difficulty in urination; also referred to as benign prostatic hypertrophy.

**Desmoid tumor:** A type of benign, locally invasive fibrous tumor capable of growing anywhere in the body.

**Essential tremor (ET):** A chronic, incurable condition with unknown cause characterized by involuntary, rhythmic tremor of a body part, most typically the hands and arms.

**High intensity focused ultrasound (HIFU):** A surgical noninvasive procedure that uses focused high energy sound waves to destroy target tissues in the body.

**Leiomyoma:** A benign tumor that can be found in the uterus, commonly called a fibroid.

**Menorrhagia:** Excessive uterine bleeding occurring at the expected intervals of the menstrual periods.

**MRI (magnetic resonance imaging):** The use of a nuclear magnetic resonance spectrometer to produce electronic images of specific atoms and molecular structures in solids, especially human cells, tissues, and organs.

**Myomectomy:** A surgical procedure to remove only fibroids; is frequently the chosen treatment for premenopausal women who want to bear more children, because it usually can preserve fertility.

**Uterine Fibroids:** Benign fibrous tissue collected in the uterine wall.

**Ventralis intermediate nucleus of the thalamus (Vim):** A part of the brain involved with movement.

**Coding**

*The following codes for treatments and procedures applicable to this document are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.*

**When services may be Medically Necessary when criteria are met:**

This Medical Policy provides assistance in understanding Healthy Blue's standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications****CPT**

0398T Magnetic resonance image guided high intensity focused ultrasound (MRgFUS), stereotactic ablation lesion, intracranial for movement disorder including stereotactic navigation and frame placement when performed

**ICD-10 Diagnosis**

G25.0 Essential tremor

**When services are Investigational and Not Medically Necessary:**

For the procedure code listed above when criteria are not met and for all other non-oncologic diagnoses.

**When services are Investigational and Not Medically Necessary:****CPT**

0071T Focused ultrasound ablation of uterine leiomyomata, including MR guidance; total leiomyomata volume less than 200 cc of tissue

0072T Focused ultrasound ablation of uterine leiomyomata, including MR guidance; total leiomyomata volume greater or equal to 200 cc of tissue

**ICD-10 Diagnosis**

All diagnoses

**When services are also Investigational and Not Medically Necessary:**

When the code describes a procedure indicated in the Position Statement section as investigational and not medically necessary.

**CPT**

55899 Unlisted procedure, male genital system [when specified as image-guided focused ultrasound ablation of prostate tissue for non-oncologic indications, such as benign prostatic hyperplasia]

60699 Unlisted procedure, endocrine system [when specified as image-guided focused ultrasound ablation for non-oncologic indications]

**HCPCS**

C9734 Focused ultrasound ablation/therapeutic intervention, other than uterine leiomyomata, with magnetic resonance (MR) guidance

**ICD-10 Diagnosis**

All non-oncologic diagnoses

**References****Peer Reviewed Publications:**

This Medical Policy provides assistance in understanding Healthy Blue's standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

---

**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications**

---

1. Altinel Y, Alkhalfan F, Qiao N, Velimirovic M. Outcomes in lesion surgery versus deep brain stimulation in patients with tremor: A systematic review and meta-analysis. *World Neurosurg.* 2019; 123:443-452.e8.
2. Avedian RS, Bitton R, Gold G, et al. Is MR-guided high-intensity focused ultrasound a feasible treatment modality for desmoid tumors? *Clin Orthop Relat Res.* 2016; 474(3):697-704.
3. Bachmann G. Expanding treatment options for women with symptomatic uterine leiomyomas: timely medical breakthroughs. *Fertil Steril.* 2006; 85(1):46-47.
4. Bond AE, Shah BB, Huss DS, et al. Safety and efficacy of focused ultrasound thalamotomy for patients with medication-refractory, tremor-dominant Parkinson disease: a randomized clinical trial. *JAMA Neurol.* 2017; 74(12):1412-1418.
5. Chang JW, Park CK, Lipsman N, et al. A prospective trial of magnetic resonance-guided focused ultrasound thalamotomy for essential tremor: Results at the 2-year follow-up. *Ann Neurol.* 2018; 83(1):107-114.
6. Chung SR, Baek JH, Suh CH, et al. Efficacy and safety of high-intensity focused ultrasound (HIFU) for treating benign thyroid nodules: a systematic review and meta-analysis. *Acta Radiol.* 2020; 61(12):1636-1643.
7. Clark NA, Mumford SL, Segars JH. Reproductive impact of MRI-guided focused ultrasound surgery for fibroids: a systematic review of the evidence. *Curr Opin Obstet Gynecol.* 2014; 26(3):151-161.
8. Cosgrove GR, Lipsman N, Lozano AM, et al. Magnetic resonance imaging-guided focused ultrasound thalamotomy for essential tremor: 5-year follow-up results. *J Neurosurg.* 2022; 138(4):1028-1033.
9. Elias WJ, Huss D, Voss T, et al. A pilot study of focused ultrasound thalamotomy for essential tremor. *N Engl J Med.* 2013; 369(7):640-648.
10. Elias WJ, Lipsman N, Ondo WG, et al. A randomized trial of focused ultrasound thalamotomy for essential tremor. *N Engl J Med.* 2016; 375(8):730-739.
11. Elble R, Comella C, Fahn S, et al. Reliability of a new scale for essential tremor. *Mov Disord.* 2012; 27(12):1567-1569.
12. Fennessy FM, Tempany CM, McDannold NJ, et al. Uterine leiomyomas: MR imaging-guided focused ultrasound surgery: results of different treatment protocols. *Radiology.* 2007; 243(3):885-893.
13. Ferreira JJ, Mestre TA, Lyons KE, et al; MDS Task Force on Tremor and the MDS Evidence Based Medicine Committee. MDS evidence-based review of treatments for essential tremor. *Mov Disord.* 2019; 34(7):950-958.
14. Friebe RW, Lin HC, Hinh PP, et al. The impact of minimally invasive surgeries for the treatment of symptomatic benign prostatic hyperplasia on male sexual function: a systematic review. *Asian J Androl.* 2010; 12(4):500-508.
15. Giordano M, Caccavella VM, Zaed I, et al. Comparison between deep brain stimulation and magnetic resonance-guided focused ultrasound in the treatment of essential tremor: a systematic review and pooled analysis of functional outcomes. *J Neurol Neurosurg Psychiatry.* 2020; 91(12):1270-1278.
16. Gizzo S, Saccardi C, Patrelli TS, et al. Magnetic resonance-guided focused ultrasound myomectomy: safety, efficacy, subsequent fertility and quality-of-life improvements, a systematic review. *Reprod Sci.* 2014; 21(4):465-476.
17. Halpern CH, Santini V, Lipsman N, et al. Three-year follow-up of prospective trial of focused ultrasound thalamotomy for essential tremor. *Neurology.* 2019; 93(24):e2284-e2293.
18. Hrary M, Segar DJ, Hayes MT, Cosgrove GR. Unilateral thalamic deep brain stimulation versus focused ultrasound thalamotomy for essential tremor. *World Neurosurg.* 2019; 126:e144-e152.
19. Iacopino DG, Gagliardo C, Giugno A, et al. Preliminary experience with a transcranial magnetic resonance-guided focused ultrasound surgery system integrated with a 1.5-T MRI unit in a series of patients with essential tremor and Parkinson's disease. *Neurosurg Focus.* 2018; 44(2):E7.

---

This Medical Policy provides assistance in understanding Healthy Blue's standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications**

---

20. Ikink ME, Voogt MJ, Verkooijen HM, et al. Mid-term clinical efficacy of a volumetric magnetic resonance-guided high-intensity focused ultrasound technique for treatment of symptomatic uterine fibroids. *Eur Radiol*. 2013; 23(11):3054-3061.
21. Jeanmonod D, Werner B, Morel A, et al. Transcranial magnetic resonance imaging-guided focused ultrasound: noninvasive central lateral thalamotomy for chronic neuropathic pain. *Neurosurg Focus*. 2012; 32(1):E1.
22. Ji Y, Hu K, Zhang Y, et al. High-intensity focused ultrasound (HIFU) treatment for uterine fibroids: a meta-analysis. *Arch Gynecol Obstet*. 2017; 296(6):1181-1188.
23. Jung HH, Chang WS, Rachmilevitch I, et al. Different magnetic resonance imaging patterns after transcranial magnetic resonance-guided focused ultrasound of the ventral intermediate nucleus of the thalamus and anterior limb of the internal capsule in patients with essential tremor or obsessive-compulsive disorder. *J Neurosurg*. 2015; 122(1):162-168.
24. Kim HS, Baik JH, Pham LD, Jacobs MA. MR-guided high-intensity focused ultrasound treatment for symptomatic uterine leiomyomata: long-term outcomes. *Acad Radiol*. 2011; 18(8):970-976.
25. Kovatcheva R, Vlahov J, Stoinov J, et al. US-guided high-intensity focused ultrasound as a promising non-invasive method for treatment of primary hyperparathyroidism. *Eur Radiol*. 2014; 24(9):2052-2058.
26. Krishna V, Fishman PS, Eisenberg HM, et al. Trial of globus pallidus focused ultrasound ablation in Parkinson's disease. *N Engl J Med*. 2023; 388(8):683-693.
27. Krishna V, Sammartino F, Agrawal P, et al. Prospective tractography-based targeting for improved safety of focused ultrasound thalamotomy. *Neurosurgery*. 2019; 84(1):160-168.
28. Lennon JC, Hassan I. Magnetic resonance-guided focused ultrasound for Parkinson's disease since ExAblate, 2016-2019: a systematic review. *Neurol Sci*. 2021; 42(2):553-563.
29. Leslie TA, Kennedy JE. High-intensity focused ultrasound principles, current uses, and potential for the future. *Ultrasound Q*. 2006; 22(4):263-272.
30. Lipsman N, Schwartz ML, Huang Y, et al. MR-guided focused ultrasound thalamotomy for essential tremor: a proof-of-concept study. *Lancet Neurol*. 2013; 12(5):462-468.
31. Louis ED. Treatment of medically refractory essential tremor. *N Engl J Med*. 2016; 375(8):792-793.
32. Madersbacher S, Schatzl G, Djavan B, et al. Long-term outcome of transrectal high-intensity focused ultrasound therapy for benign prostatic hyperplasia. *Eur Urol*. 2000; 37(6):687-694.
33. Martin E, Jeanmonod D, Morel A, et al. High-intensity focused ultrasound for noninvasive functional neurosurgery. *Ann Neurol*. 2009; 66(6):858-861.
34. Meng Y, Solomon B, Boutet A, et al. Magnetic resonance-guided focused ultrasound thalamotomy for treatment of essential tremor: A 2-year outcome study. *Mov Disord*. 2018; 33(10):1647-1650.
35. Miller WK, Becker KN, Caras AJ, et al. Magnetic resonance-guided focused ultrasound treatment for essential tremor shows sustained efficacy: a meta-analysis. *Neurosurg Rev*. 2022; 45(1):533-544.
36. Monteith S, Snell J, Eames M, et al. Transcranial magnetic resonance-guided focused ultrasound for temporal lobe epilepsy: a laboratory feasibility study. *J Neurosurg*. 2016; 125(8):1557-1564.
37. Park YS, Jung NY, Na YC, Chang JW. Four-year follow-up results of magnetic resonance-guided focused ultrasound thalamotomy for essential tremor. *Mov Disord*. 2019; 34(5):727-734.
38. Rabinovici J, Inbar Y, Revel A, et al. Clinical improvement and shrinkage of uterine fibroids after thermal ablation by magnetic resonance-guided focused ultrasound surgery. *Obstet Gynecol*. 2007; 30(5):771-777.
39. Roehrborn CG. Benign prostatic hyperplasia: an overview. *Rev Urol*. 2005; 7 Suppl 9:S3-S14.
40. Schatzl G, Madersbacher S, Djavan B, et al. Two-year results of transurethral resection of the prostate versus four 'less invasive' treatment options. *Eur Urol*. 2000; 37(6):695-701.

---

This Medical Policy provides assistance in understanding Healthy Blue's standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

---

**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications**

---

41. Schrag A. Focused ultrasound ablation of the globus pallidus internus for Parkinson's disease. *N Engl J Med*. 2023; 388(8):759-760.
42. Schreglmann SR, Krauss JK, Chang JW, et al. Functional lesional neurosurgery for tremor: a systematic review and meta-analysis. *J Neurol Neurosurg Psychiatry*. 2018; 89(7):717-726.
43. Sinai A, Nassar M, Eran A, et al. Magnetic resonance-guided focused ultrasound thalamotomy for essential tremor: a 5-year single-center experience. *J Neurosurg*. 2019: 1-8.
44. Stacy MA, Elble RJ, Ondo WG, et al; TRS study group. Assessment of interrater and intrarater reliability of the Fahn-Tolosa-Marin Tremor Rating Scale in essential tremor. *Mov Disord*. 2007; 22(6):833-838.
45. Stewart EA, Gedroyc WM, Tempany CM, et al. Focused ultrasound treatment of uterine fibroid tumors: safety and feasibility of a noninvasive thermoablative technique. *Am J Obstet Gynecol*. 2003; 189(1):48-54.
46. Stewart EA, Gostout B, Rabinovici J, et al. Sustained relief of leiomyoma symptoms by using focused ultrasound surgery. *Obstet Gynecol*. 2007; 110(2 Pt 1):279-287.
47. Stewart EA, Rabinovici J, Tempany CM, et al. Clinical outcomes of focused ultrasound surgery for the treatment of uterine fibroids. *Fertil Steril*. 2006; 85(1):22-29.
48. Taran FA, Tempany CM, Regan L, et al. Magnetic resonance-guided focused ultrasound (MRgFUS) compared with abdominal hysterectomy for treatment of uterine leiomyomas. *Ultrasound Obstet Gynecol*. 2009; 34(5):572-578.
49. Welton T, Cardoso F, Carr JA, et al. Essential tremor. *Nat Rev Dis Primers*. 2021; 7(1):83.
50. Yuen J, Miller KJ, Klassen BT, et al. Hyperostosis in combination with low skull density ratio: A potential contraindication for magnetic resonance imaging-guided focused ultrasound thalamotomy. *Mayo Clin Proc Innov Qual Outcomes*. 2021; 6(1):10-15.

**Government Agency, Medical Society and Other Authoritative Publications:**

1. American College of Obstetricians and Gynecologists. ACOG Practice Bulletin No. 228. Management of Symptomatic Uterine Leiomyomas. 2021.
2. Burke CT, Funaki BS, Ray CE. American College of Radiology (ACR) Appropriateness Criteria® treatment of uterine leiomyomas. Revised in 2017. Available at: <https://acsearch.acr.org/docs/69508/Narrative/>. Accessed on July 22, 2023.
3. Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD). Other Treatments for Fibroids. Last Reviewed: November 2, 2018. Available at: <http://www.nichd.nih.gov/health/topics/uterine/conditioninfo/treatments/Pages/radiological.aspx>. Accessed on July 13, 2023.
4. Health Quality Ontario. Magnetic Resonance-Guided Focused Ultrasound Neurosurgery for Essential Tremor: A Health Technology Assessment. *Ont Health Technol Assess Ser*. 2018; 18(4):1-141.
5. Pouratian N, Baltuch G, Elias WJ, Gross R. American Society for Stereotactic and Functional Neurosurgery Position Statement on Magnetic Resonance-Guided Focused Ultrasound for the Management of Essential Tremor. *Neurosurgery*. 2020; 87(2):E126-E129.
6. Pron G. Magnetic resonance-guided high-intensity focused ultrasound (MRgHIFU) treatment of symptomatic uterine fibroids: an evidence-based analysis. *Ont Health Technol Assess Ser*. 2015; 15(4):1-86.
7. U.S. Food and Drug Administration (FDA) Center for Devices and Radiologic Health (CDRH) 510 (k) Premarket Notification Database. InSightec Exablate Neuro. Summary of Safety and Effectiveness. No. P150038. Rockville, MD: FDA. July 11, 2016. Available at [http://www.accessdata.fda.gov/cdrh\\_docs/pdf15/P150038B.pdf](http://www.accessdata.fda.gov/cdrh_docs/pdf15/P150038B.pdf). Accessed on July 13, 2023.

---

This Medical Policy provides assistance in understanding Healthy Blue's standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue's Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications**

**Websites for Additional Information**

1. National Institute of Health (NIH): National Institute of Neurological Disorders and Stroke (NINDS): NINDS Essential Tremor Information Page. Last reviewed on June 7, 2023. Available at: <https://www.ninds.nih.gov/Disorders/All-Disorders/Essential-Tremor-Information-Page>. Accessed on July 13, 2023.
2. National Institute of Health (NIH): National Institute of Diabetes and Digestive and Kidney Diseases. Prostate Enlargement: Benign Prostatic Hyperplasia. September 2014. Available at: <https://www.niddk.nih.gov/health-information/urologic-diseases/prostate-problems/prostate-enlargement-benign-prostatic-hyperplasia>. Accessed on July 13, 2023.
3. U.S. National Library of Medicine. Uterine fibroids. Available at: <http://www.nlm.nih.gov/medlineplus/ency/article/000914.htm>. Accessed on July 13, 2023.

**Index**

Estrogen Therapy  
 ExAblate 2000 System  
 ExAblate Neuro  
 Fibroids  
 Myomectomy  
 Thalamotomy

**The use of specific product names is illustrative only. It is not intended to be a recommendation of one product over another, and is not intended to represent a complete listing of all products available.**

**Document History**

Status	Date	Action
Reviewed	08/10/2023	Medical Policy & Technology Assessment Committee (MPTAC) review. Updated Rationale, Background and References sections.
Revised	08/11/2022	MPTAC review. Added medically necessary criteria for essential tremor. Updated Coding, Rationale, Background and References sections.
Reviewed	02/17/2022	MPTAC review. Updated Description, Rationale and References sections.
Reviewed	02/11/2021	MPTAC review. Updated Rationale and References sections. Updated Coding section, added NOC 60699.
	12/16/2020	Updated Coding section with 01/01/2021 HCPCS changes; code C9747 deleted 12/31/2020.
Reviewed	02/20/2020	MPTAC review. Updated Rationale, References and Websites sections.
Reviewed	03/21/2019	MPTAC review. Updated Description and References sections.
Reviewed	09/13/2018	MPTAC review. Updated Rationale, References and Websites sections.
Reviewed	11/02/2017	MPTAC review. The document header wording updated from “Current Effective Date” to “Publish Date.”
	07/01/2017	Updated Coding section with 07/01/2017 HCPCS changes.

This Medical Policy provides assistance in understanding Healthy Blue’s standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue’s Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.



**MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications**

Revised	02/02/2017	MPTAC review. Revised investigational and not medically necessary position statement indications to include essential tremor. Updated Description, Rationale, Background, Definitions, References and Websites sections.
Revised	11/03/2016	MPTAC review. Title changed from <i>MRI Guided High Intensity Focused Ultrasound Ablation of Uterine Fibroids</i> to <i>MRI Guided High Intensity Focused Ultrasound Ablation for Non-Oncologic Indications</i> . Revised investigational and not medically necessary position statement to include all non-oncologic indications. Updated Description, Rationale, Background, Definitions, Coding, References and Websites sections.
Reviewed	05/05/2016	MPTAC review. Updated Description, Rationale, Background and References sections. Removed ICD-9 codes from Coding section.
Reviewed	05/07/2015	MPTAC review. Updated Rationale and References sections.
Reviewed	05/15/2014	MPTAC review. Updated Rationale and References sections.
Reviewed	05/09/2013	MPTAC review.
Reviewed	05/10/2012	MPTAC review. Rationale and References sections updated.
Reviewed	05/19/2011	MPTAC review. Rationale and References sections updated.
Reviewed	05/13/2010	MPTAC review. Rationale and References sections updated.
Reviewed	05/21/2009	MPTAC review. Rationale and References sections updated.
Reviewed	05/15/2008	MPTAC review. Rationale, Background and References sections updated.
	02/21/2008	The phrase "investigational/not medically necessary" was clarified to read "investigational and not medically necessary." This change was approved at the November 29, 2007 MPTAC meeting.
Reviewed	05/17/2007	MPTAC review. References updated.
Reviewed	06/08/2006	MPTAC review. Rationale, Background, Definitions and References sections updated. Removed CMS NCD which was added November 2005 in error. Position statement unchanged.
	11/21/2005	Added reference for Centers for Medicare and Medicaid Services (CMS) – National Coverage Determination (NCD).
Revised	07/14/2005	MPTAC review. Revision based on Pre-merger Anthem and Pre-merger WellPoint Harmonization.

<b>Pre-Merger Organizations</b>	<b>Last Review Date</b>	<b>Document Number</b>	<b>Title</b>
Anthem, Inc.		No prior document	
WellPoint Health Networks, Inc.	06/24/2004	2.09.18	MRI Guided High Intensity Focused Ultrasound Ablation of Uterine Fibroids

This Medical Policy provides assistance in understanding Healthy Blue’s standard Medicaid benefit plan. When evaluating coverage for a specific member benefit, reference to federal and state law, as well as contractual requirements may be necessary, since these may differ from our standard benefit plan. In the event of a conflict with standard plan benefits, federal, state and/or contractual requirements will govern. Before using this policy, please check all federal, state and/or contractual requirements applicable to the specific benefit plan coverage. Healthy Blue reserves the right to modify its Policies and Guidelines as necessary and in accordance with legal and contractual requirements. This Medical Policy is provided for informational purposes. It does not constitute medical advice. Healthy Blue may also use tools and criteria developed by third parties, to assist us in administering health benefits. Healthy Blue’s Policies and Guidelines are intended to be used in accordance with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.